

Physical and chemical structure of the solar type protostar NGC1333-IRAS4

S. Maret (sebastien.maret@cesr.fr)

*Centre d'Etude Spatiale des Rayonnements, CESR/CNRS-UPS, BP 4346, F-31028
- Toulouse cedex 04*

C. Ceccarelli

Observatoire de Bordeaux, BP 89, F-33270 Floirac

E. Caux

*Centre d'Etude Spatiale des Rayonnements, CESR/CNRS-UPS, BP 4346, F-31028
- Toulouse cedex 04*

NGC1333-IRAS4 is a binary protostellar system (IRAS4 A and B) located in the south part of the Perseus cloud. They have been classified as Class 0 protostars (Andre et al., 2000) and are associated with molecular outflows probed by CO and CS millimeter lines.

We observed IRAS4 with the *Long Wavelength Spectrometer* (Clegg et al., 1996) on board the *Infrared Space Observatory* (Kessler et al., 1996) in grating mode. These observations consist of a spectral survey of the central position containing both IRAS4A and B, and two positions along the CO outflow (see also Ceccarelli et al., 1999). We detected fourteen H₂O lines, nine CO lines, and the [OI] and [CII] fine structure lines. No significant water and CO emission was found along the outflow, whereas the [OI] and [CII] lines were detected with the same intensity than on the central position.

The observed molecular emission can a priori have three different origins: the outflows, a photodissociation region (PDR), and the collapsing envelope around the protostar. The [OI] and [CII] emission can be relatively well explained by a PDR with a density from 10^4 to 10^6 cm⁻³, and an incident FUV field, whose flux is of the order of the average interstellar FUV field flux. On the contrary, both CO and H₂O emission could arise either in shocked gas (e.g. due to the impact of the outflowing gas with the envelope) and/or in the envelope itself.

In the past we (Ceccarelli et al., 1996) computed the thermal emission of a protostellar envelope. This model computes in a self consistent way the thermal balance, chemistry and radiative transfer in the envelope, in the “inside-out” framework (Shu, 1977). We successfully applied this model to IRAS4, and we were able to explain the H₂O emission as due to an envelope collapsing towards an object whose mass is $0.6 M_{\odot}$ and accreting at $5.10^{-5} M_{\odot} \text{ yr}^{-1}$ (see Maret et al., in preparation, for more details). Assuming a constant mass accretion rate this gives an age of



© 2008 Kluwer Academic Publishers. Printed in the Netherlands.

12000 yr. The water abundance is about 10^{-6} with respect to H_2 in the outer cold envelope, and it is enhanced by one order of magnitude in the central region, where the grain mantles evaporate. The existence of such *hot-core* like region has been also claimed in the case of the solar type protostar IRAS16293-2422, where a similar study was carried out (Ceccarelli et al., 2000a; Ceccarelli et al., 2000b). IRAS16293-2422 seems to be more massive ($0.8 M_{\odot}$) than IRAS4 and accreting at a lower accretion rate ($3.10^{-5} M_{\odot} \cdot \text{yr}^{-1}$), suggesting that IRAS16293-2422 is more evolved than IRAS4, with an estimated age of 27000 yr. This conclusion is in agreement with the relatively large millimeter continuum observed in IRAS4, which implies a large amount of dust surrounding this source. Finally, both the hot core like region and the region in which CO-rich ices are predicted to evaporate (i.e. when the dust temperature is ~ 25 K) are larger in IRAS16293-2422 than in IRAS4, which can explain the larger CO depletion observed in the latter (Blake et al., 1995).

This study also emphasizes the necessity of ground based observations, where higher spatial and spectral resolutions are achievable. H_2CO and CH_3OH are of particular interest as they are among the most abundant components of the grain mantles, and are therefore expected to evaporates in the innermost parts of the envelope. Appropriate transitions can hence be used to constrain the physical and chemical conditions in such innermost parts of protostellar envelopes.

References

- Andre, P., D. Ward-Thompson, and M. Barsony: 2000, 'From Prestellar Cores to Protostars: the Initial Conditions of Star Formation'. *Protostars and Planets IV* pp. 59+.
- Blake, G. A., G. Sandell, E. F. van Dishoeck, T. D. Groesbeck, L. G. Mundy, and C. Aspin: 1995, 'A molecular line study of NGC 1333/IRAS 4'. *ApJ* **441**, 689–701.
- Ceccarelli, C., A. Castets, E. Caux, D. Hollenbach, L. Loinard, S. Molinari, and A. G. G. M. Tielens: 2000a, 'The structure of the collapsing envelope around the low-mass protostar IRAS 16293-2422'. *A&A* **355**, 1129–1137.
- Ceccarelli, C., D. J. Hollenbach, and A. G. G. M. Tielens: 1996, 'Far-Infrared Line Emission from Collapsing Protostellar Envelopes'. *ApJ* **471**, 400+.
- Ceccarelli, C., L. Loinard, A. Castets, A. G. G. M. Tielens, and E. Caux: 2000b, 'The hot core of the solar-type protostar IRAS 16293-2422: H_2CO emission'. *A&A* **357**, L9–L12.
- Clegg, P. E., et al.: 1996, 'The ISO Long-Wavelength Spectrometer.'. *A&A* **315**, L38–L42.
- Kessler, et al.: 1996, 'The Infrared Space Observatory (ISO) mission.'. *A&A* **315**, L27–L31.
- Shu, F. H.: 1977, 'Self-similar collapse of isothermal spheres and star formation'. *ApJ* **214**, 488–497.